

ФАРМАКОЛОГІЯ І ТОКСИКОЛОГІЯ

UDC 619:636.087.7:615.918:616.992:636

Monitoring of changes in mineral and vitamin metabolism under the influence of feed additive in piglets with experimental associated mycotoxicosis

Andriichuk A. , Melnyk A. , Vovkotrub N. 

Bila Tserkva National Agrarian University

 Andriichuk A. E-mail: andriichuk.av@gmail.com



Andriichuk A., Melnyk A., Vovkotrub N.
Monitoring of changes in mineral and vitamin metabolism under the influence of feed additive in piglets with experimental associated mycotoxicosis. *Naukovy visnyk veterinary medicine*, 2020. № 2. PP. 131–139.

Рукопис отримано: 12.11.20.

Прийнято: 27.11.20.

Затверджено до друку: 24.11.20.

doi: 10.33245/2310-4902-2020-160-2-131-139

Toxicobiological effect of mycotoxins association of the *Penicillium* and *Fusarium* fungus (T-2 toxin at a concentration of 0.1 mg/kg, fumonisin B1 – 0.5 mg/kg, vomitoxin (DON) – 0.1 mg/kg, penicillic acid – 1 mg/kg) was accompanied by the development of a complex pathological process in weaned piglets. In this regard, the detoxification and sorption capacity of the complex feed additive "Harufix+" based on mannanoligosaccharides was studied. The additive effect on the resorptive activity of mineral and vitamin nutrients of feed under the normal feeding conditions and in case of contamination with mycotoxins has been studied. The use of enterosorbent offset the toxic effects of micromycete metabolites, which contributed to the growth of piglets. Thus, weight growth rate increase of the piglets in group 1 (i.e., those whose diet included the additive, unlike the diet of the animals in control group) constituted 16 %, while their average weight growth rate was high and constituted 1.96 kg per day.

In addition, during the study of calcium, phosphorus, magnesium, ferum, zinc, copper and manganese in the piglets blood was not found excretion of these elements with a sorbent, moreover, noted the normalization their blood level.

The study content of vitamins A and E, the same as with mineral nutrients, has not been established decrease during treatment with study the pharmaceutical. The obtained results testify the active absorption in the gastrointestinal tract of the vitamin components within the fodder combined with fodder additive "Harufix+" and high biological accessibility of its transport forms. The efficiency of the additive can be explained by its composition, namely the complex of mineral and organic components that are formed by modification of the organic cations of the mineral surface.

Key words: mycotoxins, mycotoxicosis, macro- and microelements, sorbent, vitamin metabolism, piglets.

Problem statement. The health, productivity and reproductive functions of animals largely depend on the degree of contamination of feed with pathogenic microflora and toxins of various origins. Mycotoxins are fungal metabolites that may have deleterious effects in animals [1]. Mycotoxins are detected in cereal grains worldwide, with a prevalence of 88 % on feed and raw feedstuffs [2].

Mycotoxins routinely occur in common feedstuffs such as corn, corn silage, small grains and small grain silage, especially when growing conditions are sub-optimal. It should be noted that mycotoxins may or may not be present in moldy

feedstuffs. In addition, mycotoxins may be present in feeds that appear free of mold which makes mycotoxins a challenge for producers and nutritionists alike [3].

There is increasing evidence of animal feed contamination by several mycotoxins that is coming from the specialists in animal husbandry, animal feed production, veterinary medicine and mycotoxicology [4]. This can be explained by the ever growing database of toxic fungal metabolites, newly discovered facts about the synthesis of certain mycotoxins and their groups by different fungus types, the development of the system of permanent monitoring and the improvement of

the methodology of research on mycotoxins contained in animal feed.

Analysis of recent research. According to Whitlow and Hagler, mycotoxins exert their effects through several means including 1) decreased feed intake, 2) reduced nutrient absorption and impaired metabolism, 3) altered function of the endocrine and exocrine systems, 4) suppressed immune function, 5) altered intestine microbial growth and 6) changes in white blood cell and neutrophil counts [5].

Müller et al. conducted an experimental investigation of weaner pigs and the influence of combined administration of ochratoxin A, fumonisin B₁, deoxynivalenol and T2 toxin in quantities expected to be present in feeds of central European origin. They observed toxic effect of mycotoxins that slightly surpassed the toxico-biological effect after single administration of ochratoxin A [6]. The contamination of animal feed with several toxic compounds of micromycetes (i.e., micro-fungi) occurs quite frequently, since many kinds of *Aspergillus*, *Penicillium* and *Fusarium* produce more than one mycotoxin, and various fungi may contaminate grain components of animal compound feed. Thus, corn and wheat naturally contaminated with vomitoxin (i.e., deoxynivalenol, DON), 15-acetyl DON, Fusarium acid (FA) and zearalenone were included in the diet of start pigs.

Feeding pigs with purified mycotoxins fumonisin B₁ and aflatoxin B₁ together and separately influenced the immune system, biochemical, hematological and clinical parameters [7]. With the joint administration of mycotoxins, these indicators were more pronounced, and the toxic effects intensified and sometimes very significantly, especially liver disease syndrome [8].

The study demonstrated considerable decrease in growth rate, feed consumption and brain neurochemistry alterations [9]. The combined contamination of feedstuff with fungal toxins complicates the prevention of mycotoxicoses in animals, because mycotoxins have a wide range of physico-chemical properties. Thus, application of only one method of detoxication and decontamination (e.g., the use of specific enterosorbent) is not always efficient [10].

Besides, it is well known that sorbents tend to bind (adsorb) and withdraw from the organism macro- and microelements, vitamins, and nutrients, which subsequently leads to the decrease in animal productivity and becomes the reason of the rejection of mycotoxin-binding pharmaceuticals.

Schell et al. used weanling and growing pigs to investigate the effect of sodium bentonite clay (1 %) on mineral metabolism in diets with or without aflatoxin. Feeding aflatoxin contaminated

feed increased phosphorus (P), sodium (Na) and Zn absorption and retention suggesting a possible increased metabolic demand for these minerals when aflatoxin is present.

In addition, feeding sodium bentonite decreased Mg absorption regardless of the presence of aflatoxin. The addition of bentonite clay also decreased calcium (Ca) and Na absorption and retention in aflatoxin contaminated diets and decreased Na absorption when the feed was free of aflatoxin. Similar to the effects of Hydrated sodium calcium aluminosilicate, Zn absorption and retention was decreased in diets supplemented with sodium bentonite. The effects of feeding sodium bentonite clay on iron (Fe) was confounded in this study by the increase in dietary Fe (446 ppm vs. 292 ppm) from the addition of the clay [11].

In addition to negatively affecting Zn, bentonite has also been shown to decrease copper (Cu) bioavailability in sheep [12] fed no supplemental trace minerals. Although the cation composition was not disclosed, bentonite was fed at 0.5% of the diet (as fed basis). In this study, bentonite decreased the ruminal solubility of Zn, Cu and Mg and led to significant decreases in Cu in both plasma (0.75 vs. 0.71 µg/ml) and liver (602 vs. 504 µg/g DM).

In conclusion, it appears utilizing silicate minerals as mycotoxin sequestering agents could lead to decreases in both Zn and Cu status. The interaction between complexed trace minerals and mycotoxin sequestering agents has not been researched. However, providing a portion of supplemental Zn and Cu as amino acid complexes may be warranted to improve the likelihood of maintaining optimal trace mineral status when diets contain silicate-based mycotoxin sequestering agents [13].

Other mycotoxin sequestering agents include activated charcoal, cholestyramine, chlorophyllin and yeast cell wall-derived agents. Although these may be beneficial at reducing the impact of mycotoxins in humans, aquatic and other animal species, there is currently no data available on their interaction with mineral nutrition [14].

The aim of the study was to analyze the changes in vitamin and mineral metabolism in piglets under the influence of the feed additive Harufix+ in associated mycotoxicosis.

Material and methods. For the purposes of this study we formed 4 groups of weaner pigs, 10 piglets in each. The piglets in group 1 were administered with combined feed that contained Harufix+ in dosage of 1 kilo per ton of feed. The piglets in group 2 were fed with the feed that contained T₂ toxin (0.1 mg/kg), fumonisin B₁ (0.5 mg/kg), vomitoxin (deoxynivalenol, 0.1 mg/kg) and penicillic acid (1 mg/kg). In order to produce T₂ toxin we used *Fusarium sporotrichiella*, strain 2M

which was produced at the State poultry research station of National Academy of Agrarian sciences of Ukraine). Other mycotoxins by micromycete isolates were incubated at the Department of microbiology and virology of Bila Tserkva National Agrarian University. The diet of group 3 piglets included a complex of mycotoxins and anti-toxic feed additive "Harufix+" (1 kg/ton). The feed of the animals of group 4 did not contain mycotoxins (control group). The experiment lasted 14 days. At the beginning and during the experimental period the animals were weighed. At the end of experimental period it was conducted biochemical laboratory analysis of piglets' blood samples for evaluation the values of calcium, phosphorus, magnesium, zinc, manganese, ferum, copper, vitamins A and E. Blood for the study was taken from the piglets' orbital venous sinus in vacuum tubes with gel and coagulation activator. The blood serum content of total calcium was determined in the reaction with calcium arsenase III, inorganic phosphorus – by UV-detection of phosphomolybdate complex, total magnesium – with the calmagite indicator, vitamin A – by the method of Bessey in the modification of VI Levchenko, vitamin E – in reaction with 2,2-dipyridyl. All these methods were carried out with reagents of research and production association "Philisit-diagnostics" using a semi-automatic biochemical analyzer Stat Fax (USA). The serum content of ferum, copper, zinc and manganese was determined by atomic absorption spectrophotometry using an atomic absorption spectrophotometer Shimadzu (Japan).

Statistical processing of the results was performed using Statistica 10 (StatSoft Inc., USA, 2011).

Results. Based on animals' weight tests, we come to conclusion that fodder additive "Harufix+" has positive impact on body weight growth. Thus, weight growth rate increase of the piglets in group 1 (i.e., those whose diet included the pharmaceutical, unlike the diet of the animals in control group) constituted 16 %, while their average weight growth rate was high and constituted 1.96 kg per day. This observation provides indirect evidence in support of the conclusion that this fodder additive does not bind and withdraws from the organism nutrients, vitamins, macro- and microelements.

We compared weight growth data of animals in group 2 (whose feed was contaminated with mycotoxins) and group 3 (whose diet was contaminated as well, but also included the feed additive). In comparison, we discovered the difference in 5 %. Average weight growth rate in group 2 was low and constituted 2.05 kg per day, which was the effect of the mycotoxins in the feed. The same parameter in group 3 constituted 2.15 kg per day (Figure 1). The difference in growth rates in these groups is apparent and strongly testifies to the protective capacity of Harufix+ in induced mycotoxicosis in weaner pigs. In the first group, the rate of weight gain was the highest and averaged 2.73 kg, which is due to the efficient use and absorption of feed nutrients against the background of additional use of feed additives.

The analysis of mineral and vitamins status in piglets did not reveal any disruption of their homeostasis. Furthermore, we observed the normalization of the ratio of certain elements of their mineral nutrition.

The content of total calcium in group 1 remained comparable to those in control group and




Fig. 1. Weight growth changes in piglets.

it was 2.51 ± 0.14 mmol/L (2.23 – 2.67 mmol/L). It should be noted that in piglets of other experimental groups, the content of total calcium in the blood also did not have a significant difference compared with controls and animals of the first group (Table 1). The content of non-organic phosphorus in blood serum of piglets in group 1 was significantly higher ($p < 0.001$; $+18.2\%$) relatively the corresponding parameter in group 4 (2.75 ± 0.055) and it was in average 3.25 ± 0.020 mmol/L (3.21 – 3.28). Also it should be noted the significantly increasing in 1.2 times ($p < 0.001$) the phosphorus content in animals of group 3, which received a complex of mycotoxins and anti-toxic additive “Harufix+” relatively the control group.

The changes in serum magnesium homeostasis were characterized by a probable decrease in its blood level in animals of the second group on average to 0.73 ± 0.047 mmol/L ($p < 0.05$) compared with the control and 3rd groups, which may be due to impaired absorption of this macronutrient in the intestine or reduction of its reabsorption in the renal tubules due to the complex action of mycotoxins on these organs. In other experimental groups of piglets, the total magnesium serum content in the probably did not differ from the control (Table 1).

Ferum is a microelement that is in particular demand in the organisms of young, fast growing animals. Thus, Ferum was abundant in the blood

of animals, whose diet included Harufix+, its content averaged 681.7 ± 151.9 $\mu\text{g}/100$ ml. The Ferum blood content of the piglets of groups 3 and 4 almost did not differ and averaged 505.8 ± 182.1 (207.0 – 835.5) and 529.0 ± 268.0 (212.0 – 1063.7) $\mu\text{g}/100$ ml respectively. Accordingly, in animals whose diet contained mycotoxins its content was on a considerably lower level and averaged 384.2 ± 178.0 $\mu\text{g}/100$ ml, which is 1.8 times less than in group 1 and 1.4 times less than in control group (Table 2).

Copper metabolism in the organism of those piglets that were administered this fodder additive did not undergo significant changes. The content of this micronutrient in animals of group 1 averaged 288.9 ± 13.8 (261.7 – 306.5) as opposed to 285.9 ± 42.8 $\mu\text{g}/100$ ml in those in control (Table 2). However, in the second experimental group, the blood copper content in piglets was 31.7% lower compared to the control group, which indicates a negative effect of mycotoxins on the absorption of this trace element in the gastrointestinal tract of animals.

The serum Zinc content in group 1 averaged 38.7 ± 2.31 $\mu\text{g}/100$ ml, which is 34.8% higher ($p < 0.01$) than in control group (28.7 ± 1.85). The highest level was in piglets of 3rd experimental group – 70.9 ± 31.1 $\mu\text{g}/100$ ml, which is 2.2 and 2.5 times higher than the same index in groups 2 and 4, respectively (Table 2). Probably, this is due to the

Table 1 – The changes in macronutrient metabolism indexes in piglets

Indicator	Group			
	1	2	3	4
Ca total, mmol/L	2.23–2.67 2.51 ± 0.14	2.31–2.75 2.53 ± 0.127	2.50–2.88 2.68 ± 0.11	2.44–2.53 2.49 ± 0.027
P non-organic, mmol/L	3.21–3.28 $3.25 \pm 0.020^{***}$	2.74–3.36 2.95 ± 0.203	3.18–3.35 $3.26 \pm 0.050^{***}$	2.5–2.84 2.75 ± 0.055
Mg total, mmol/L	0.95–1.14 1.02 ± 0.06	0.64–0.80 $0.73 \pm 0.047^*$	1.05–1.2 1.12 ± 0.04	0.74–1.33 1.11 ± 0.184

Note: * $p < 0.05$; *** $p < 0.001$ relatively control (group 4).

Table 2 – The changes in micronutrient metabolism indexes in piglets

Indicator	Group			
	1	2	3	4
Fe, $\mu\text{g}/100$ ml	397.9–917.7 681.7 ± 151.9	203.8–740.2 384.2 ± 178.0	207.0–835.5 505.8 ± 182.1	212.0–1063.7 529.0 ± 268.0
Cu, $\mu\text{g}/100$ ml	261.7–306.5 288.9 ± 13.8	54.7–265.6 195.2 ± 70.2	246.4–268.5 260.7 ± 7.14	230.3–370.1 285.9 ± 42.8
Zn, $\mu\text{g}/100$ ml	34.2–41.8 $38.7 \pm 2.31^{**}$	28.2–36.9 31.5 ± 2.73	37.5–133.1 70.9 ± 31.1	26.7–32.4 28.7 ± 1.85
Mn, $\mu\text{g}/100$ ml	19.6–34.0 $25.9 \pm 4.25^*$	13.0–46.8 26.1 ± 10.5	16.0–39.2 26.0 ± 6.88	15.2–16.6 15.9 ± 0.40

Note: * $p < 0.05$; ** $p < 0.01$ relatively control (group 4).

positive effect of Harufix+ components on the zinc absorption against the background of mycotoxins damage.

Regarding changes in manganese levels, in piglets of all three experimental groups, its content in the blood averaged almost at the same level – 25.9–26.1 $\mu\text{g}/100\text{ ml}$, which is 1.6 times higher than the average value in animals of the control group ($15.9 \pm 0.40\text{ }\mu\text{g}/100\text{ ml}$). However, a probable increase in blood manganese levels was only in group 1 ($p < 0.05$).

Similarly to the microelements values, our analysis of the vitamins A and E levels did not reveal their decrease that would result from the pharmaceutical use. Thus, the value of vitamin A in blood serum of group 1 was 14.3–25.8 and averaged $18.6 \pm 3.64\text{ }\mu\text{g}/100\text{ ml}$, whereas the normal indicator for the piglets of this age is 20–50 $\mu\text{g}/100\text{ ml}$ (Figure 2).

The highest average value of vitamin A was in the piglets blood of the control group – $29.1 \pm 4.6\text{ }\mu\text{g}/100\text{ ml}$, while the lowest content was observed in animals of the 2nd experimental group, which was 1.9 times less than the control group ($p < 0.05$). The analysis of blood serum for the value of vitamin A in group 3 was 38 % higher then in group 2 and 13.4 % – then in group 1 (Figure 2).

It should be noted that changes in the content of vitamin E in the blood of piglets under the complex influence of the association of mycotoxins without the use of enterosorbent were characterized by a significant decrease to $0.17 \pm 0.02\text{ mg}/100\text{ ml}$ ($p < 0.05$) compared with animals of the first and third experimental groups. However, low level of vitamin E was observed in piglets of the control group – $0.19 \pm 0.03\text{ mg}/100\text{ ml}$, which also did not take this feed additive (Figure 3).




Fig. 2. The blood vitamins A content in piglets.




Fig. 3. The blood vitamins E content in piglets.

The content of vitamin E in 67% of animals of groups 2 and 4 was less than the minimum level of 0.2 mg/100 ml. While in groups 1 and 3 we can observe a positive effect of the feed additive “Harufix+” components on the absorption and stability of tocopherol homeostasis, because the average value in the blood of piglets in these groups was in the range of 0.25–0.26 mg/100 ml.

Discussion. Minerals provide an important role in animals. Their importance is in maintaining a certain osmotic pressure of blood plasma, acid-base balance, permeability of various biological membranes, regulation of enzyme activity, preservation of the structures of biomolecules, maintenance of the motor and secretory functions of the digestive tract.

We found that in case of mycotoxicosis caused by toxins of *Fusarium* and *Penicillium* fungus, there were significant deviations in the content of total magnesium in the blood plasma of piglets, the level of which was 34, 2% less than in the control group. The use of the sorption additive “Harufix+” did not have a negative effect on the absorption of this macronutrient in the intestine, since its content in the blood of piglets of the 1st and 3rd experimental groups did not significantly differ from the control group.

The positive effect of feed additives on absorption and the absence of its sorption effect on phosphorus in the gastrointestinal tract of pigs was evidenced by a probable increase in its blood level in animals of groups 1 and 3 compared with controls ($p < 0,001$). Phosphorus absorption in pigs is limited, depends on the sufficient amounts of vitamin D₃ and the presence of calcium in their nutrition [15, 16]. Apparently, such active transport of this macroelement is associated with the sorptive qualities of the Harufix+ components related to phosphorus accumulation and its transport into blood vessels. The significance of non-organic phosphates in the organism is determined not only by the fact that their great amounts are concentrated in bone tissue in the compound of calcium phosphate. Monosodium and disodium phosphates establish phosphorus buffer system in blood, which along with carbonate and protein buffers participates in acid-alkaline balance regulation. In parallel, phosphorus plays an important role in the processes of phosphorylation and dephosphorylation. This provides kidney absorption and excretion, as well as lipids and proteins transport [17]. This is demonstrated by the piglets' body mass increase of group 1, i.e. in those that were administered fodder additive within their combined nutrition.

However, the blood total calcium content in piglets of all experimental groups remained at the

level of control and was within physiological limits. That is, neither the use of feed additives nor the complex effect of the association of mycotoxins did not cause deviations in the homeostasis of this macronutrient. Probably, this can be explained by a rather severe complex and multicomponent endogenous system of ensuring the stability of calcium blood value, which involves mainly humoral factors and vitamin D also. Ensuring the calcium homeostasis stability and maintaining its physiological blood plasma level is carried out by various mechanisms, the main of which are changes in the degree of intestine absorption, the proximal renal tubules reabsorption and the calcium's mobilization from the bone component [18].

The impact of Harufix+ on micronutrients absorption reveals itself in its positive influence on Zinc transport, which was reflected in a probable increase its blood level in piglets of the first group by 34.8% compared to control ($p < 0,01$). The best result regarding Zinc homeostasis was obtained when using the sorbent in piglets with the associated influence of mycotoxins – 70.9 ± 31.1 µg/100 ml, which is 2.5 times higher than the average value in the control group. Biochemical function of Zinc in the organism is related to the activity of ferments, which need Zinc as a necessary component or an activator. Up to date, Zinc was found present in more than 200 metalloenzymes, which take part in a range of metabolic processes, including synthesis and decomposition of carbohydrates, fats, proteins and nucleic acids [18].

It is important to sustain microelements homeostasis on sufficiently stable level since it determines their synergism in relation to metabolism stimulation in general. For instance, Ferum is highly needed for normal activity of dehydrogenases, catalase and peroxidase; Copper is needed for oxygenase, xanthine oxidase and urate oxidase; Manganese – for transferases; Magnesium – for phosphohydrolase and Zinc for carbonic anhydrase and carboxypeptidase [19, 20]. It is important to note that physiological anemia is the typical outcome of the insufficient Ferum in piglets organisms during the first weeks of their lives. Anemia causes the death of 20 to 30% suckling piglets during the first weeks of their lives. Our research has shown positive dynamics in the assimilation of Ferum in piglets that consumed a feed additive, both as part of common feed and against the background of its contamination by mycotoxins. In the second experimental group, the Ferum's serum content was the lowest – 384.2 ± 178.0 µg/100 ml.

The use of Harufix+ did not adversely affect the absorption of Copper and Manganese in the intestines of piglets, which was reflected in the constant levels of these micronutrients in the blood of ani-

mals of groups 1 and 3. Moreover, in the blood of piglets that consumed the feed additive, there was a probable increase in 1.6 times the content of serum Manganese compared to the control group ($p < 0,05$).

The study content of vitamins A and E, the same as with mineral nutrients, has not been established decrease during treatment with study the feed additive. The obtained results testify to the active absorption in the gastrointestinal tract of the vitamin components within the fodder combined with fodder additive "Harufix+" and high biological accessibility of its transport forms. The efficiency of the pharmaceutical can be explained by its composition. Thus, the complex of mineral and organic components is formed by means of modification of organic cations of the mineral surface. As a result, not merely a mixture of organic and mineral components is formed, but a new organic complex. Acarbose that is a component of Harufix+ in combination the mineral components absorb mycotoxins and excrete them from the animal's organism, as well as normalize the microflora of the intestines. Beta-glucan, which is also contained in this fodder additive, improves the function of the gastrointestinal tract, activating the enzymatic system of its mucous membrane.

Conclusions. 1. The study demonstrated anti-toxic efficiency of fodder additive "Harufix+" in pigs with mycotoxicosis experimentally induced by a complex of mycotoxins, which was manifested by its efficient sorption properties characteristics relative to T-2 toxin, fumonisin B1, vomitoxin and penicillic acid.

2. This fodder additive did not cause the disruption of beneficial components of nutrition digestion, it proved biologically harmless, and positively impacted body weight increase in the animals.

3. Monitoring of changes in the content of total calcium, inorganic phosphorus, magnesium, iron, zinc, copper, manganese and vitamins A and E in the piglets blood confirmed the stability of their homeostasis against the background of the sorbent.

Information on compliance with ethical standards. Experimental studies were conducted in compliance with the requirements of the Law of Ukraine No. 3447 – IV of 21.02.06 "On the animals protection from cruel treatment" and in accordance with the basic principles of the "European Convention for the protection of vertebrate animals used for experimental and scientific purposes" (Strasbourg, 1986), the Declaration on the Humane Treatment of Animals (Helsinki, 2000) and the National Congress on Bioethics "General Ethical Principles of Animal Experiments" (Kyiv, 2001).

Information about the interest conflict. The authors declare no interest conflict.

LIST OF LITERATURE

1. Abdallah M.F., Girgin G., Baydar T. Occurrence, prevention, and limitation of mycotoxins in feeds. *Anim. Nutr. Feed Technol.* 2015. 15. P. 471–490.
2. Gruber-Dorninger C., Jenkins T., Schatzmayr G. Global mycotoxin occurrence in feed: A ten-year survey. *Toxins.* 2019. 11. 375 p.
3. Висланько О.О., Зінов'єв С.Г., Гиря В.М. Ефективність використання нового сорбенту мікотоксинів у свинарстві. *Вісник Полтавської державної аграрної академії.* 2010. № 2. С. 107–110.
4. Paterson R.R., Lima N. *Toxicology of Mycotoxins. Molecular, Clinical and Environmental Toxicology.* 2010. Vol. 100. P. 31–63.
5. Whitlow L.W., Hagler W.M. Mold and mycotoxin issues in dairy cattle: Effects, prevention and treatment. *Proc. Southeast Dairy Herd Mgmt Conf., Macon, GA.* 2008. P. 80–89.
6. Mueller G., Kielstein H., Rosner A. Studies of the influence of combined administration of ochratoxin A, fumonisin B1, deoxynivalenol and T-2 toxin on immune and defence reactions in weaner pigs. *Mycoses.* 1999. Vol. 42. P. 485–493.
7. Effects of dietary fumonisin Br-containing culture material, deoxynivalenol contaminated wheat, or their combination on growing barrows/ R.B. Harvey et al. *Am. J. Vet. Res.* 1996. №57. P. 1790–1794.
8. The effects of mycotoxins, fumonisin B1 and aflatoxin B1, on primary swine alveolar macrophages / B.H. Liu et al. *Toxicol. Appl. Pharmacol.* 2002. Vol. 180. P. 197–204.
9. Swamy H.V.L.N., Smith T.K., MacDonald H.J. Effect of feeding blends of grains naturally contaminated with *Fusarium* mycotoxins on brain regional neurochemistry of starter pigs and broiler chickens. *Anim. Sci.* 2004. Vol. 82. P. 2131–2139.
10. Діаз Д. Микотоксини и микотоксикозы. М.: Печатный город, 2006. 382 с.
11. Schell T.C., Lindemann M.D., Kornegay E.T., Blodgett D.J. Effects of feeding aflatoxin-contaminated diets with and without clay to weanling and growing pigs on performance, liver function, and mineral metabolism. *J. Anim. Sci.* 1993. 71. P. 1209–1218.
12. Ivan M., Dayrell D.S., Hidiogrou M. Effects of bentonite and monensin on selected elements in the stomach and liver of fauna-free and faunated sheep. *J. Dairy Sci.* 1992. 75. P. 201–208.
13. Effects of hydrated sodium calcium aluminosilicate on fescue toxicosis and mineral absorption/ A.B. Chestnut et al. *J. Anim. Sci.* 1992.70. P. 2838–2846.
14. CAST, Council for Agricultural Science and Technology. *Mycotoxins: Risks in plant, animal and human systems. Task Force Report.* Ames, IA. 2003. no. 139.
15. Федак Н.М., Вовк Я.С., Чумаченко С.П., Душара И.В. Минеральные вещества в кормлении сельскохозяйственных животных. Предгорное и горное земледелие и животноводство. 2012. № 54 (1). С. 128–135.
16. Дозозалежний вплив вітаміну Е на обмін холекальциферолу в організмі /Л.І. Апуховська та ін. *Вісник Білоцерків. держ. аграр. ун-ту. Вип. 29. Біла Церква,* 2004. С. 3–15.
17. Кондрахін І.П. Метаболічний синдром: сучасне уявлення, перспективи використання. *Біологія тварин.* Львів, 2010. Т. 12 (№ 2). С. 63–66.

18. Ветеринарна клінічна біохімія: підручник / Левченко В.І. та ін.; за ред. В.В. Влізла. Біла Церква: БНАУ, 2019. 416 с.

19. Марченко Ф.С., Сторожук Т.В. Хелатные микроэлементы – важный компонент комбикормов и премиксов. Зерновые продукты и комбикорма. 2010. № 1. С. 37–38.

20. Камышников В.С. Справочник по клинико-биохимическим исследованиям и лабораторной диагностике. М.: МЕДпресс-информ, 2004. 920 с.

REFERENCES

1. Abdallah, M.F., Girgin, G., Baydar, T. (2015). Occurrence, prevention, and limitation of mycotoxins in feeds. Anim. Nutr. Feed Technol. 15, pp. 471–490.

2. Gruber-Dorninger, C., Jenkins, T., Schatzmayr, G. (2019). Global mycotoxin occurrence in feed: A ten-year survey. Toxins. 11, 375 p.

3. Vyslanko, O.O., Zinoviev, S.H., Hyria, V.M. (2010). Efektyvnist vykorystannia novoho sorbentu mikotoksyniv u svynarstvi [Efficiency of using a new sorbent of mycotoxins in pig breeding]. Visnyk Poltavskoi derzhavnoi ahraryoi akademii [Bulletin of the Poltava State Agrarian Academy]. no. 2, pp. 107–110.

4. Paterson, R.R., Lima, N. (2010). Toxicology of Mycotoxins. Molecular, Clinical and Environmental Toxicology. Vol. 100, pp. 31–63.

5. Whitlow, L.W., Hagler, W.M. (2008). Mold and mycotoxin issues in dairy cattle: Effects, prevention and treatment. Southeast Dairy Herd Mgmt Conf., Macon, GA. Jr. pp 80–89.

6. Mueller, G., Kielstein, H., Rosner, A. (1999). Studies of the influence of combined administration of ochratoxin A, fumonisin B1, deoxynivalenol and T-2 toxin on immune and defence reactions in weaner pigs. Mycoses. Vol. 42, pp. 485–493.

7. Harvey, R.B., Edrington, T.S., Kubena, L.F. (1996). Effects of dietary fumonisin B1-containing culture material, deoxynivalenol contaminated wheat, or their combination on growing barrows. Am. J. Vet. Res. no. 57, pp. 1790–1794.

8. Liu, B.H., Yu, F.Y., Chan, M.H. (2002). The effects of mycotoxins, fumonisin B1 and aflatoxin B1, on primary swine alveolar macrophage. Toxicol. Appl. Pharmacol. Vol. 180, pp. 197–204.

9. Swamy, H.V.L.N., Smith, T.K., MacDonald, H.J. (2004). Effect of feeding blends of grains naturally contaminated with Fusarium mycotoxins on brain regional neurochemistry of starter pigs and broiler chickens. Anim. Sci. Vol. 82, pp. 2131–2139.

10. Dyaz, D. (2006). Mikotoksiny i mikotoksikozy [Mycotoxins and mycotoxicoses]. М.: Printing town, 382 p.

11. Schell, T.C., Lindemann, M.D., Kornegay, E.T., Blodgett, D.J. (1993). Effects of feeding aflatoxin-contaminated diets with and without clay to weanling and growing pigs on performance, liver function, and mineral metabolism. J. Anim. Sci. 71, pp. 1209–1218.

12. Ivan, M., Dayrell, D.S., Hidirolou, M. (1992). Effects of bentonite and monensin on selected elements in the stomach and liver of fauna-free and faunated sheep. J. Dairy Sci. 75, pp. 201–208.

13. Chestnut, A.B., Anderson, P.D., Cochran, M.A., Frébourg, H.A., Gwinn, K.D. (1992). Effects of hydrated sodi-

um calcium aluminosilicate on fescue toxicosis and mineral absorption. J. Anim. Sci. 70, pp. 2838–2846.

14. CAST, Council for Agricultural Science and Technology. (2003). Mycotoxins: Risks in plant, animal and human systems. Task Force Report. Ames, IA. no. 139.

15. Fedak, N.M., Vovk, Ya.S., Chumachenko, S.P., Dushara, Y.V. (2012). Mineral'nye veshchestva v kormlenii sel'skohozjajstvennykh zhivotnykh [Minerals in the feeding of farm animals]. Predgornoe i gornoe zemledelie i zhivotnovodstvo [Foothill and mountain agriculture and animal husbandry]. no. 54(1), pp. 128–135.

16. Apukhovska, L.I., Nikiforova, T.M., Romanova, S.A. (2004). Dozozaleznyi vplyv vitaminu E na obmin kholekaltsyferolu v orhanyzmi [Dose-dependent effect of vitamin E on the metabolism of cholecalciferol in the body]. Visnyk Bilotserkiv. derzh. ahrary. un-tu. [Bulletin of Bila Tserkva State Agrarian University]. Bila Tserkva, Issue 29, pp. 3–15.

17. Kondrakhin, I.P. (2010). Metabolichnyi syndrom: suchasne uiavlennia, perspektyvy vykorystannia [Metabolic syndrome: modern idea, prospects of use]. Biolohiia tvaryn [Animal biology]. Lviv, Vol. 12 (№ 2), pp. 63–66.

18. Levchenko, V.I., Vlizlo, V.V., Kondrakhin, I.P. (2019). Vetrynarna klinichna biokhimiia: pidruchnyk [Veterinary clinical biochemistry: a textbook]. Bila Tserkva: BNAU, 416 p.

19. Marchenkov, F.S., Storozhuk, T.V. (2010). Helatnye mikrojelementy – vazhnyj komponent kombikormov i premiksiv [Chelated trace elements are an important component of compound feed and premixes]. Zernovye produkty i kombikorma [Cereals and compound feed]. no. 1. pp. 37–38.

20. Kamysnikov, V.S. (2004). Spravochnik po kliniko-biokhymicheskim issledovaniyam i laboratornoj diagnostike [Reference book on clinical and biochemical research and laboratory diagnostics]. М.: MEDpress-inform, 920 p.

Моніторинг зміни показників мінерально-вітамінного метаболізму під впливом кормової добавки за експериментального асоційованого мікотоксикозу в поросят

Андрійчук А.В., Мельник А.Ю., Вовкотруб Н.В.

Токсикобіологічна дія асоціації мікотоксинів роду *Penicillium* і *Fusarium* (Т-2 токсин у концентрації 0,1 мг/кг, фумонізін В1 – 0,5 мг/кг, вомітоксин (ДОН) – 0,1 мг/кг, пеніцилова кислота – 1 мг/кг) супроводжувалася розвитком комплексного патологічного процесу в організмі відлучених поросят. У зв'язку із цим було досліджено дезінтоксикаційну та сорбційну здатність комплексної кормової добавки "Харуфікс+" на основі маннанолігосахаридів. Вивчено вплив сорбенту на резорбтивну активність мінеральних і вітамінних нутрієнтів корму за звичайних умов годівлі та в разі контамінації мікотоксинами. Застосування ентеросорбенту нівелювало токсичну дію метаболітів мікроміцетів, що сприяло зростанню приростів поросят. Так, різниця приростів поросят першої групи, які отримували з кормом добавку, порівняно з тваринами контрольної групи, становила 16 %, причому середньодобовий приріст тварин першої групи був найвищим і дорівнював 1,96 кг.

Водночас під час дослідження вмісту кальцію, фосфору, магнію, феруму, цинку, купруму та мангану в крові поросят не було встановлено виведення цих елемен-

тів із сорбентом, більше того, відзначали нормалізацію їх рівня.

За результатами дослідження вмісту вітамінів А і Е, аналогічно як і з мінеральними нутрієнтами, не було встановлено їх зниження на фоні застосування досліджуваного сорбенту. Отримані результати свідчать про активне всмоктування у шлунково-кишковому тракті вітамінних компонентів у складі корму в поєднанні з кормовою добавкою "Харуфікс+" та високу біологічну доступність його транспортних форм. Така ефективність добавки пояснюється, насамперед, її складом, а саме комплексом мінеральних і органічних компонентів, які формуються шляхом модифікації органічними катіонами поверхні мінералу.

Ключові слова: мікотоксини, мікотоксикоз, макро- і мікроелементи, сорбент, вітамінний обмін, поросята.

Мониторинг изменения показателей минерально-витаминного метаболизма под влиянием кормовой добавки при экспериментальном ассоциированном микотоксикозе у поросят

Андрейчук А.В., Мельник А.Ю., Вовкотруб Н.В.

Токсикобиологическое действие ассоциации микотоксинов рода *Penicillium* и *Fusarium* (Т-2 токсин в концентрации 0,1 мг/кг, фумонизин В1 – 0,5 мг/кг, vomitоксин (ДОН) – 0,1 мг/кг, пеницилловая кислота – 1 мг/кг) сопровождалось развитием комплексного патологического процесса в организме отлученных поросят. В связи с этим было изучено дезинтоксикационную и сорбционную способность комплексной кормовой добавки "Харуфикс+" на основе маннанолигосахаридов. Изучено влия-

ние сорбента на резорбтивную активность минеральных и витаминных нутриентов корма в обычных условиях кормления и в случае контаминации микотоксинами. Применение энтеросорбента нивелировало токсическое действие метаболитов микромицетов, что способствовало росту приростов поросят. Так, разница приростов поросят первой группы, получавших с кормом добавку, по сравнению с животными контрольной группы, составляла 16 %, причем среднесуточный прирост животных первой группы был высоким и составлял 1,96 кг.

Наряду с этим в ходе исследования содержания кальция, фосфора, магния, железа, цинка, меди и марганца в крови поросят не было установлено связывания этих элементов с сорбентом, более того, отмечали нормализацию их уровня в крови.

По результатам исследования содержания витаминов А и Е, аналогично как и с минеральными нутриентами, не было установлено их снижения на фоне применения испытуемого сорбента. Полученные результаты свидетельствуют об активном всасывании в желудочно-кишечном тракте витаминных компонентов в составе корма в сочетании с кормовой добавкой "Харуфикс+" и высокую биологическую доступность его транспортных форм. Такая эффективность добавки объясняется, прежде всего, ее составом, а именно комплексом минеральных и органических компонентов, которые формируются путем модификации органическими катионами поверхности минерала.

Ключевые слова: микотоксины, микотоксикозы, макро- и микроэлементы, сорбент, витаминный обмен, поросята.



Copyright: © Andriichuk A., Melnyk A., Vovkotrub N. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.



Andriichuk A.
Melnyk A.
Vovkotrub N.

ID <https://orcid.org/0000-0001-9144-5272>
ID <https://orcid.org/0000-0001-9129-4814>
ID <https://orcid.org/0000-0003-3297-454X>